

# Current Biology

## Three Phases of Ancient Migration Shaped the Ancestry of Human Populations in Vanuatu

### Highlights

- New ancient DNA supports a shift in ancestry during early migrations to Vanuatu
- A single spread from New Britain can explain most of the ancestry of later groups
- More recent Polynesian migrations contributed both cultural and genetic legacies

### Authors

Mark Lipson, Matthew Spriggs, Frederique Valentin, ..., Nadin Rohland, Ron Pinhasi, David Reich

### Correspondence

mlipson@hms.harvard.edu (M.L.), Matthew.Spriggs@anu.edu.au (M.S.), ron.pinhasi@univie.ac.at (R.P.), reich@genetics.med.harvard.edu (D.R.)

### In Brief

Lipson et al. report new genetic data and analyses shedding light on three human migrations to Vanuatu. The first involved people with primarily East Asian-related ancestry; the second, shortly afterward, and likely following a similar route from New Britain, primarily Papuan ancestry; and the third, more recently, Polynesian ancestry.



## Article

# Three Phases of Ancient Migration Shaped the Ancestry of Human Populations in Vanuatu

Mark Lipson,<sup>1,2,\*</sup> Matthew Spriggs,<sup>3,4,\*</sup> Frederique Valentin,<sup>5</sup> Stuart Bedford,<sup>4,6,7</sup> Richard Shing,<sup>4</sup> Wanda Zinger,<sup>8</sup> Hallie Buckley,<sup>9</sup> Fiona Petchey,<sup>10,11</sup> Richard Matanik,<sup>12</sup> Olivia Cheronet,<sup>13</sup> Nadin Rohland,<sup>1</sup> Ron Pinhasi,<sup>13,\*</sup> and David Reich<sup>1,2,14,15,16,\*</sup>

<sup>1</sup>Department of Genetics, Harvard Medical School, Boston, MA 02115, USA

<sup>2</sup>Department of Human Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA

<sup>3</sup>School of Archaeology and Anthropology, College of Arts and Social Sciences, The Australian National University, Canberra, ACT 2601, Australia

<sup>4</sup>Vanuatu National Museum, Vanuatu Cultural Centre, Port Vila, Vanuatu

<sup>5</sup>MSH Mondes, CNRS, UMR 7041, 92023 Nanterre, France

<sup>6</sup>Department of Archaeology and Natural History, College of Asia-Pacific, The Australian National University, Canberra, ACT 2601, Australia

<sup>7</sup>Max Planck Institute for the Science of Human History, 07745 Jena, Germany

<sup>8</sup>Muséum national d'Histoire naturelle, UMR 7194 (HNHP), MNHN/CNRS/UPVD, Sorbonne Université, Musée de l'Homme, 75016 Paris, France

<sup>9</sup>Department of Anatomy, School of Biomedical Sciences, University of Otago, Dunedin 9054, New Zealand

<sup>10</sup>Radiocarbon Dating Laboratory, Division of Health, Engineering, Computing and Science, University of Waikato, Hamilton 3240, New Zealand

<sup>11</sup>ARC Centre of Excellence for Australian Biodiversity and Heritage, College of Arts, Society and Education, James Cook University, Cairns, QLD 4878, Australia

<sup>12</sup>Lelema World Heritage Committee and Vanuatu Cultural Centre, Port Vila, Vanuatu

<sup>13</sup>Department of Evolutionary Anthropology, University of Vienna, 1090 Vienna, Austria

<sup>14</sup>Medical and Population Genetics Program, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

<sup>15</sup>Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115, USA

<sup>16</sup>Lead Contact

\*Correspondence: [mlipson@hms.harvard.edu](mailto:mlipson@hms.harvard.edu) (M.L.), [Matthew.Spriggs@anu.edu.au](mailto:Matthew.Spriggs@anu.edu.au) (M.S.), [ron.pinhasi@univie.ac.at](mailto:ron.pinhasi@univie.ac.at) (R.P.), [reich@genetics.med.harvard.edu](mailto:reich@genetics.med.harvard.edu) (D.R.)

<https://doi.org/10.1016/j.cub.2020.09.035>

## SUMMARY

The archipelago of Vanuatu has been at the crossroads of human population movements in the Pacific for the past three millennia. To help address several open questions regarding the history of these movements, we generated genome-wide data for 11 ancient individuals from the island of Efate dating from its earliest settlement to the recent past, including five associated with the Chief Roi Mata's Domain World Heritage Area, and analyzed them in conjunction with 34 published ancient individuals from Vanuatu and elsewhere in Oceania, as well as present-day populations. Our results outline three distinct periods of population transformations. First, the four earliest individuals, from the Lapita-period site of Teouma, are concordant with eight previously described Lapita-associated individuals from Vanuatu and Tonga in having almost all of their ancestry from a "First Remote Oceanian" source related to East and Southeast Asians. Second, both the Papuan ancestry predominating in Vanuatu for the past 2,500 years and the smaller component of Papuan ancestry found in Polynesians can be modeled as deriving from a single source most likely originating in New Britain, suggesting that the movement of people carrying this ancestry to Remote Oceania closely followed that of the First Remote Oceanians in time and space. Third, the Chief Roi Mata's Domain individuals descend from a mixture of Vanuatu- and Polynesian-derived ancestry and are related to Polynesian-influenced communities today in central, but not southern, Vanuatu, demonstrating Polynesian genetic input in multiple groups with independent histories.

## INTRODUCTION

A key distinction within Pacific studies has been between Near Oceania, the part of the Western Pacific (comprising New Guinea; the Bismarck Archipelago, including New Britain and New Ireland; and the main Solomon Islands) settled for approximately 50,000 years by modern humans, and

Remote Oceania [1]. Remote Oceania encompasses the whole of Micronesia and Polynesia and the geographically designated Melanesian island groups of Vanuatu, New Caledonia, and Fiji (as well as the scattered islands of the Reefs and Santa Cruz groups in the southeast Solomons), which were only settled starting around 3,000 years before present (BP) [1].

Vanuatu is a key archipelago in the history of Pacific settlement given its status both as the first major island group in southern Remote Oceania to be occupied by humans and as an important regional crossroads during the succeeding three millennia [2, 3]. Our understanding of the genetic history of Vanuatu has been advanced by three studies reporting genome-wide ancient DNA data from individuals who lived in the archipelago over the course of its human settlement [4–6]. The earliest sampled individuals, who belong to the first human migration to Vanuatu (labeled by some commentators as Migration 1 or M1 [7, 8]), are associated with early phases of the Lapita cultural complex and likely with the initial spread of Austronesian languages into Oceania (where Austronesian is now by far the most widespread language family) [9, 10]. They had almost entirely East Asian-related ancestry, from a source that originated in Taiwan and has been termed “First Remote Oceanian” (FRO) [4]. Later individuals (including present-day people, who identify as “Ni-Vanuatu”), by contrast, have largely Papuan ancestry likely originating in New Britain, which reached the Reefs-Santa Cruz [11] and Vanuatu [5, 6] either during latest Lapita or post-Lapita times after 2800 BP. (We use the term “Papuan” to refer to the deep ancestral lineage that contributes the majority of the ancestry found in present-day populations from Near Oceania.) Previous papers differed in their interpretation of this second migration (M2) as being either a time-constrained event [6] or a slower process of continuing genetic exchange through time [5].

Previous studies [5, 6] also noted but did not address in detail signals of a third distinct migration stream (M3) occurring within the last millennium and associated with the establishment of “Polynesian Outlier” communities in Vanuatu (as in other areas of Melanesia and Micronesia): that is, islands where Polynesian sub-group languages are spoken and where elements of Polynesian material and non-material culture are practiced [12, 13]. Polynesian impacts in Vanuatu also extend to a number of islands neighboring the Outlier communities showing Polynesian influence but without full language replacement. Little is known, however, about the degree of population movement accompanying these Polynesian-derived cultural and linguistic changes [14, 15].

One such Polynesian-influenced island is Efate in central Vanuatu, where two Polynesian-language-speaking communities exist today, one on the small off-shore island of Ifira and one at Mele on the southwest of the island. Also located on Efate and the adjacent small islands of Eretok and Lelepa is “Chief Roi Mata’s Domain,” which was inscribed on the UNESCO World Heritage Area list in 2008 on the basis of strong links between oral traditions and a spectacular mortuary site excavated in the 1960s [16]. Some versions of the local oral traditions and aspects of the associated material culture have suggested strong Polynesian influence, illustrated by stories about Chief Roi Mata and his political role on Efate and adjacent islands of the Shepherd Group [16, 17]. The burial site at Eretok was thought initially to date to the 13<sup>th</sup> century CE [16], but subsequent radiocarbon dates from Eretok and from Mangaas (Mangaasi), the village site on Efate said to have been the home of Chief Roi Mata and his closest followers [16], now place the burials at c. 1600 CE [18].

To gain a genetic perspective on the history of Chief Roi Mata’s Domain, and more generally on the history of Polynesian influence in Vanuatu, we sampled three individuals from the

Eretok (also known as Retoka or “Hat Island”) Island complex where Roi Mata was buried (according to tradition) along with two individuals from sub-floor burials at Mangaas for ancient DNA analysis. We also report new genome-wide ancient DNA data from six additional individuals from Efate, complementing published data [4, 6]: four from the Teouma Lapita cemetery (~3000–2750 BP, thus doubling the sample size available from that site), one from the Taplins 1 rockshelter, and one from Banana Bay. We combined these 11 individuals with 26 ancient Vanuatu individuals from the literature (who have previously not been analyzed together) [4–6], eight other published ancient Oceanian individuals, and diverse present-day populations to shed light on the following primary questions pertaining to the population movements referred to above as M1, M2, and M3:

M1. Does the increased sample of Lapita-period burials from Teouma, combined with other sites, reveal a more diverse founding population than was previously documented?

M2. Can we better elucidate the source, timing, and duration of Papuan migration into Vanuatu?

M3. Do the newly reported individuals from Eretok and Mangaas within the Chief Roi Mata’s Domain World Heritage Area show particular relatedness to Polynesians as some oral traditions and features of the archaeological record would suggest?


## RESULTS

### Sample and Data Preparation

We generated genome-wide ancient DNA data for 11 new individuals (Figure 1; Table 1; STAR Methods; Data S1A) and increased sequencing coverage for one previously reported individual from Teouma [6] (I5951/TeoQE, previously 23,107 sites covered, now 120,830). In dedicated clean rooms, we extracted DNA from either petrous bone samples (Teouma, Mangaas, and two Eretok individuals) or teeth (Taplins, one Eretok, and Banana Bay) and prepared next-generation sequencing libraries, enriching for a set of ~1.2 million single-nucleotide polymorphisms (SNPs). Based on a combination of criteria, all yielded authentic ancient DNA (STAR Methods). We created genotype data for analysis by assigning the observed base from one randomly chosen sequencing read covering each targeted SNP. For most analyses, we merged the new data with published data from both ancient and present-day Oceanians [4–6] (Data S1B). We also obtained three new radiocarbon dates to help establish chronology in relation to previously dated samples [19]; notably, the dates from Eretok and Mangaas confirm that the individuals lived within the past several centuries (Table 1; Data S1C).

### PCA

We began by performing a principal component analysis (PCA) in which we computed axes by using Kankanaey (Philippines), Naisioi (Solomon Islands), and New Guinea Highlanders and projected all other individuals (STAR Methods; Figure 2). Visually, PC1 corresponds to relative proportions of FRO ancestry (lower on the left, higher on the right), whereas PC2 corresponds to affinity to populations from the Solomon Islands versus New Guinea (up and down, respectively). Present-day groups from



**Figure 1. Geographic Context**

(A) Regional map.

(B) Map of Vanuatu.

(C) Map of Efate with sample sizes for newly reported individuals from each site.

**Table 1. Information for Newly Reported Individuals**

Lab ID	Skeletal Code (Element)	Date	Location	Sex	Mt Hap	Y Hap	SNPs
I5265	Teo_B10A (petrous)	[3000–2750 BP]	Efate, Teouma	M	no call	O	13,594/4,469
I5266	Teo_B10C (petrous)	[3000–2750 BP]	Efate, Teouma	M	B4a1a1	O	136,137/45,599
I5267	Teo_B30B (petrous)	3170–2810 calBP (3050 ± 49 BP, Wk-22658)	Efate, Teouma	M	no call	no call	8,612/2,802
I5268	Teo_B30C (petrous)	3010–2760 calBP (2995 ± 21 BP, Wk-22659)	Efate, Teouma	M	no call	no call	4,165/1,396
I6188	TAP_E149 (tooth)	[2600–2200 BP]	Efate, Mele-Taplins	M	Q1b	C1b2a	23,812/8,088
I10966	Musée de l'Homme 25788 (petrous)	[500–200 BP]	Efate, Mangaas	F	Q1	n/a	648,879/230,929
I10967	Musée de l'Homme 25787 (petrous)	290–0 calBP (180 ± 20 BP, PSUAMS-5494)	Efate, Mangaas	F	Q2a3	n/a	469,594/167,469
I10968	Musée de l'Homme 25793 (petrous)	[500–200 BP]	Eretok	M	B4a1a1	C1b2a	848,415/295,552
I10969	Musée de l'Homme 25791 (petrous)	[500–200 BP]	Eretok	F	P2	n/a	749,208/267,632
I14493	Musée de l'Homme 25797 (tooth)	490–310 calBP (350 ± 20 BP, PSUAMS-6698)	Eretok	M	P2	C1b2a	506,596/179,141
EFE005	EFE005 (tooth)	310–0 calBP (234 ± 19 BP, MAMS-29695)	Efate, Banana Bay	M	P1d2	C1b2a	74,434/25,228

Date, calibrated radiocarbon date (95.4% CI) or burial context estimate (brackets); Mt/Y hap, mitochondrial DNA/Y chromosome haplogroup; SNPs, unique autosomal target sites covered at least once/sites covered in primary analysis dataset. See also [Data S1A–S1C](#).

New Britain and Vanuatu form a cluster with relatively uniform values along PC2 but a moderate amount of spread along PC1, with Polynesians and Polynesian Outlier populations farther to the right. Ancient individuals mostly overlap present-day groups from the same island chains, but the Lapita-associated individuals from Teouma (Vanuatu) and Talasiu (Tonga), the ancient individuals from Malakula, and some individuals from Eretok and Mangaas fall farther to the right.

The direction of greatest variation within Vanuatu in [Figure 2](#) is approximately left to right (likely reflecting differential FRO/Papuan mixture proportions), which is well aligned with the primary direction of variation linking New Britain, Vanuatu, Polynesia, and the ancient Lapita-associated individuals. This pattern suggests the possibility that many or all of the populations along this extended cline can be modeled in a simple way as having a shared pair of ancestry components in different proportions: one represented by Papuan ancestry related to that found in some parts of New Britain and Vanuatu at close to 100% and one represented by FRO ancestry related to that found in the Lapita-associated individuals at close to 100% [\[4–6\]](#).

### Explicit Admixture Modeling


Guided by the PCA results, we tested candidate admixture models by using the *qpAdm* software [\[20, 21\]](#). Previous results [\[4–6, 11\]](#), as well as [Figure 2](#), indicate a high degree of regional population structure in Near Oceania, with largely distinct clusters of Papuan ancestry found in New Guinea, the Solomon Islands (excluding Santa Cruz and Polynesian Outliers), New Britain, and New Ireland, although many populations (e.g., from New Ireland) can be modeled as having mixtures of multiple Papuan ancestry components. In the following analyses, we often use Nasioi (non-Austronesian speakers from the island of Bougainville) and Baining (non-Austronesian speakers from

New Britain) to represent the Solomon Islands and New Britain clusters, respectively, because they are the populations with both the lowest proportions of FRO ancestry (~20% and ~5%) and the highest proportions of the distinctive local Papuan ancestry from their clusters in our dataset [\[4–6, 11\]](#).

For almost all of the ancient Vanuatu individuals, we obtain successful *qpAdm* models (i.e., high p values for model fit) using Baining (Marabu subgroup) and Kankanaey (Austronesian speakers from the Philippines related to the ancestors of FRO) as the two proxy sources, even with Nasioi as an outgroup ([STAR Methods](#); [Data S1D](#)). Conversely, if we use Nasioi as a proxy source in place of Baining, almost none of the models are successful. We note that poor fits can result from any unmodeled shared ancestry between the outgroups and either the test population or the proxy sources, for example from small amounts of contamination (for ancient individuals) or if the FRO-related ancestry in Nasioi (as an outgroup) is a better source than the FRO-related ancestry in Kankanaey. For Polynesians and Polynesian Outliers, our power to distinguish between different lineages is limited by their lower proportions of Papuan ancestry, but we observe similar results with better fits when using Baining rather than Nasioi as a proxy source. As previously reported [\[5\]](#), the fits improve with Malaita (a Solomon Islands population with some New Britain-related ancestry; see [Figure 2](#) and [\[6\]](#)) in place of Nasioi, but they are worse than with Baining and are rejected at  $p < 0.05$  for most populations.

The quantitative mixture proportion estimates from *qpAdm* ([Figure 3](#)) are also in good agreement with PCA. The lowest proportions of FRO ancestry we observe are 0%–3.6% and 0.6%–6.6% (truncated 95% confidence interval [CI]) for post-Lapita individuals from Efate and Tanna, respectively, and the highest proportions are 96.4%–99.2%, 96.4%–100%, and 87.4%–100% for Lapita-associated individuals from Teouma,



**Figure 2. PCA Results**

Axes were computed by using three present-day populations (bottom right legend), and other present-day (no fill) and ancient (large filled symbols; newly reported with black outline) individuals were projected and plotted by using the first two PCs. Colors correspond to genetic clusters centered around the Solomon Islands (red), New Ireland (orange), New Britain (blue), New Guinea (black), and Polynesia and Taiwan (green). N.G., New Guinea; Polyn., Polynesian; Van., Vanuatu; anc., ancient.

### Sources of Papuan and FRO Ancestry

We explored the cline of Papuan and FRO ancestry in Remote Oceania in more detail through allele-sharing symmetry tests. To allow us to compare different populations along the cline, we plotted  $f_4$  statistics of interest as a function of a


Talasiu (Tonga), and Malakula, respectively. The individuals from Chief Roi Mata's Domain are relatively variable, ranging from a low of 17.3%–22.0% total FRO ancestry for I10966 (Mangaas) to a high of 38.3%–44.2% FRO ancestry for I10969 (Eretok). We also compared ancestry proportion estimates on the autosomes and X chromosomes to test for possible sex-biased admixture. We observed isolated signals of sex bias, replicating previously reported instances for present-day Polynesians and ancient Malakula (Data S1D) [4, 5]; additional examples could exist, but our statistical power is limited by sequencing coverage and available sample sizes.

### Dates of Admixture

Previous work [4–6] has shown that the majority of present-day populations in Vanuatu have average admixture dates centered around ~2000 BP, in line with other Oceanians, although some groups, especially those with potential Polynesian-related ancestry, yield more recent dates (e.g., Futuna, ~1075 ± 225 BP [6]). We estimated dates of admixture for the Eretok and Mangaas individuals by using both MALDER [22] and DATES [23] and inferred average dates of roughly 20–30 generations, or 550–850 years, before the individuals lived (i.e., ~1400–700 BP; Table 2). This range extends somewhat earlier than the likely arrival of westward-moving Polynesian groups in Vanuatu, which, based on archaeological evidence, occurred around 1000–750 BP [13, 24]. However, under a scenario of Polynesian influx, the expected average admixture dates would reflect a combination of recent and older events, given that both Polynesians and local groups would have been admixed already. We did not detect significant evidence of multiple waves of admixture from MALDER, but because both proximal sources would have had mixtures of the same (Papuan and FRO) types of ancestry, it is difficult to disentangle the different episodes [22]. Still, the relatively recent dates for Eretok and Mangaas, together with the observed heterogeneity in mixture proportions [25], provide evidence of more recent admixture processes.

separate statistic ( $f_4(X, \text{New Guinea Highlanders}; \text{Kankanaey, Australian})$ ) proportional to FRO ancestry (Figure S1). If all test populations X can be modeled as having mixtures of ancestry related to the same two source populations (in different proportions), then such plots are expected to show a straight line (STAR Methods).

First, we computed the statistic  $f_4(X, \text{Dai}; \text{Nasioi, New Guinea Highlanders})$ , which tests for relative allele sharing between the test population X and groups from the Solomon Islands and New Guinea (Figure 4A; Data S1E). Two test populations would be expected to yield different values of this statistic (after correcting for proportions of FRO ancestry) if they have different sources for their Papuan ancestry (for example, one from New Britain and the other from New Guinea, New Ireland, or the Solomon Islands). With a few exceptions (Erromango,  $Z = -3.2$ ; Teouma,  $Z = -2.5$ ; I10969,  $Z = 2.3$ ; Tutuba,  $Z = 4.0$ ; all others within  $|Z| = 2$  of the regression line), present-day and ancient Remote Oceanians give highly uniform results (purple and green points and regression line in Figure 4A), consistent with a common source for their Papuan ancestry. Tutuba, as a copra plantation island, plausibly experienced recent admixture between Ni-Vanuatu and introduced plantation laborers from other parts of Melanesia. Why Erromango is an exception is unclear; it was a much-visited island in the 19<sup>th</sup> century by groups purchasing and cutting sandalwood and, as a result of such contacts, suffered population collapse through introduced diseases [26]. Among Near Oceanians, as expected, groups from New Guinea are generally below the Remote Oceanian line, and groups from the Solomon Islands are above. A subset of populations from New Britain, however, closely track the Remote Oceanians, suggesting that they represent good proxies for the source of Papuan ancestry that contributed (predominantly) to Vanuatu and Polynesia. We confirmed this result by using *qpWave* [27], where we obtain reasonably good two-component fits (rank 1  $p = 0.18$  without Nasioi as an outgroup,  $p = 0.02$  with Nasioi added; STAR Methods) for 10 ancient



**Figure 3. Ancestry Proportions for Ancient Vanuatu Individuals**

Results are from two-component *qpAdm* models estimating total proportions of Papuan and FRO ancestry, truncated at 0% for four individuals with negative point estimates. Newly reported individuals are represented by points with black outlines. Some points are shifted slightly left and right for legibility. Bars show two standard errors in both directions (for truncated individuals, upper limit of point estimate plus two standard errors). See Table 1 for full date intervals and Data S1D for full *qpAdm* results.

Vanuatu population groups together with present-day Tongan plus Melamela (Austronesian speakers from New Britain with  $|Z| < 2$  deviation from the regression line in Figure 4A). Present-day Vanuatu populations require four ancestry sources (rank 3  $p = 0.17$  without Nasioi as an outgroup,  $p = 0.02$  with Nasioi added), plausibly due to small proportions of distinct Papuan (as in Erromango and Tutuba) or other (e.g., East Asian or European) ancestry resulting from recent contacts.

Next, we performed similar tests for possible different sources of FRO ancestry. We first computed  $f_4(X, \text{New Guinea Highlanders}; \text{Teouma, Kankanaey})$  to test relatedness of FRO ancestry across Oceania to the Teouma individuals versus present-day Kankanaey. All populations yield positive values highly correlated with levels of FRO ancestry (Figure S2A; Data S1F), indicating that the ancestry is more closely related to the Teouma individuals [4]. We then computed  $f_4(X, \text{New Guinea Highlanders}; \text{Teouma, Talasiu})$  to test whether the FRO ancestry is more closely related to the Lapita-associated individuals from Vanuatu or from Tonga. Although our statistical power is limited by the close relationship between the two Lapita-associated groups, we obtain significantly non-zero values for populations having relatively high FRO ancestry, with the negative slope implying (slightly) greater affinity to Talasiu than to Teouma (Figure S2B; Data S1G). However, we observe only minor deviations from the regression line (max  $|Z| = 2.5$ ). Thus, the FRO ancestry found in sampled ancient and present-day Oceanian populations appears to be relatively uniform in its relationships to the Lapita-associated individuals from Vanuatu and Tonga, and slightly closer to the latter.

**Table 2. Inferred Average Dates of Admixture**

Test Group or Individual	MALDER Result (Gen/Year)	DATES Result (Gen/Year)
Eretok (triple)	16.4 ± 4.5/459 ± 126	24.1 ± 5.0/675 ± 140
Mangaas (pair)	36.5 ± 15.4/1,023 ± 432	22.6 ± 6.9/633 ± 193
I10968 (Eretok)	–	18.4 ± 4.6/514 ± 128
I10969 (Eretok)	–	29.6 ± 9.1/828 ± 256
I14493 (Eretok)	–	28.3 ± 9.3/793 ± 261
I10966 (Mangaas)	–	31.7 ± 13.5/888 ± 378
I10967 (Mangaas)	–	18.5 ± 7.4/517 ± 207


Gen/year, generations/years before the individuals lived (mean ± 1 SE).

### Polynesian Genetic Legacy

By using similar methods, we tested for the presence of specifically Polynesian-related ancestry via the statistic  $f_4(X, \text{Tolai}; \text{Kankanaey, Tongan})$  (STAR Methods; Figure 4B; Figure S3; Data S1H). As expected, other Polynesians show very strong allele sharing with Tonga ( $|Z| > 9$  for Samoa, Tahiti, and the Polynesian Outliers of Ontong Java, Rennell and Bellona, and Tikopia). Within Vanuatu, most groups are consistent with the baseline level established by Near Oceanians, but some—generally those with higher total proportions of FRO ancestry—display excess allele sharing with Tonga. These include one ~150 BP Efate (Ifira) individual ( $Z < -3$ ) and present-day Aneityum, Banks, Efate, Emae, Futuna, Makura, Mele (high-FRO subgroup, from the island of Efate), and Tongoa (all  $Z < -4$ ). Among our newly reported ancient individuals, both from Mangaas and two of the three from Eretok have strong signals of Polynesian affinity ( $-5.0 \leq Z \leq -3.6$ ).

We also attempted to determine the source of this Polynesian affinity more precisely by using statistics  $f_4(X, \text{Tolai}; \text{Polynesian1, Polynesian2})$  (Data S1I–S1L). We did not detect significant differences in allele sharing relative to Tonga versus Samoa, but for a number of Polynesian-influenced groups in Vanuatu, we observed modest excess allele sharing with Tonga versus Polynesian Outliers (max  $|Z| = 3.6, 2.5$ , and  $2.5$  for Ontong Java, Rennell and Bellona, and Tikopia, respectively). One exception was excess relatedness between Namaram (from the island of Pentecost) and Ontong Java ( $Z = 3.2$ ). However, for the most part, the source of the Polynesian-related ancestry in the Vanuatu groups appears to be slightly more closely related to populations from Polynesia than to other Polynesian Outlier communities in Melanesia (at least in their current genetic makeup).

We then tested for excess allele sharing between the Eretok and Mangaas individuals and other Vanuatu populations (STAR Methods; Data S1M–S1Q). We detected several significant signals: (1) between the five ancient individuals and present-day Efate ( $Z = 1.8$ – $3.2$ ) and especially the high-FRO subgroup of present-day Mele ( $Z = 4.2$ – $7.5$ ), (2) between the Eretok individuals I10968 and I10969 and the ~150 BP individual from Ifira ( $Z = 2.7$ – $3.6$ ), and (3) among the five Eretok and Mangaas individuals themselves ( $Z = 1.8, 2.2, 2.5, 2.7, 2.9, 3.6, 6.5, 7.2, 7.4$ , and  $16.4$ ). A separate statistic testing for allele sharing with present-day Futuna identified a strong relationship with Aneityum ( $Z > 9$ ) but confirmed no particular relatedness to Eretok or Mangaas (Data S1R). Follow-up analyses also indicated that the Eretok



**Figure 4. Allele-Sharing Regression Tests**

(A) Test for differential Papuan ancestry. The regression line was computed by using groups from Vanuatu and Polynesia, except for the Lapita-associated individuals (rightmost three points).

(B) Test for Polynesian influence. The regression line was computed by using Near Oceanian populations. Filled points represent the Eretok/Mangaas individuals. The legend is the same for both panels (the “New Guinea” label includes some closely related populations from nearby islands; some in the far lower left in (A) are omitted for scale), and bars show two standard errors in each direction. Polyn., Polynesian. See also Figures S1–S4 and Data S1E–S1R.

Britain), one of which can parsimoniously characterize the Papuan ancestry in Melamela (New Britain), Vanuatu, and Tonga.

Within Vanuatu, the model contains separate two-stage admixture histories in the southern and central parts of the archipelago. Present-day Futuna can be modeled as having 56% ancestry related to individuals from Tanna (who themselves are inferred to have 12% FRO ancestry and 88% Papuan ancestry) and 44% related to Polynesians. For Efate, I5259 (from Mangaliliu, but not necessarily associated with the Chief Roi Mata’s Domain sites) is inferred to have 11% FRO ancestry and 89% Papuan ancestry, and the Eretok/Mangaas group can be modeled as having 63% of their ancestry related to I5259 and 37% related to Polynesians (for a total of ~33% FRO ancestry). If we model Eretok/Mangaas and Futuna as having excess FRO (but not specifically Polynesian-related) ancestry, the log likelihood of the model is more than 30 units lower, with residual poorly predicted  $f$  statistics ( $Z > 5$ ). Tanna and I5259 might not be exact representatives for the true ancestral source groups,

individual I14493 and the Mangaas individual I10967 are close family relatives (probably second degree; Figure S4), explaining their especially high allele sharing ( $Z = 16.4$ ) and confirming oral traditions directly linking both sites in the Roi Mata stories.

### Admixture Graph Analysis


Finally, we built an admixture graph to explore relationships among multiple populations simultaneously, including present-day Tanna and Futuna, a ~600 BP individual (I5259) from Efate [6], Eretok and Mangaas, Polynesians, and diverse Near Oceanians (Figure 5; Figure S5; STAR Methods). The final model predicts all  $f$  statistics relating the populations to within 2.7 standard errors of their observed values. We inferred two admixture events [6] among four ancestral Papuan lineages (associated with New Guinea, the Solomon Islands, and two with New

so the inferred proportions of Polynesian-related ancestry could be slightly inaccurate, but they are plausible proxies based on both the regional genetic context and the fit quality of the final model.

### DISCUSSION

The human genetic history of Vanuatu is complex, featuring interactions between multiple populations with diverse origins. This complexity is not surprising given that the archipelago stretches for more than 1,000 km and forms a crucial intervisible link in the southwest Pacific from the Reefs and Santa Cruz (at the eastern edge of the Solomon Islands) to New Caledonia. Furthermore, in light of the great cultural diversity that characterizes Vanuatu today, it would not be surprising if different parts of





**Figure 5. Schematic of Admixture Graph Results**

Inferred phylogeny is shown for FRO-related ancestry (light green) and four Papuan lineages (pink, black, and blue, shown separately because they are not related by a simple tree). Arrows denote Papuan ancestry found in Vanuatu and Polynesia (solid blue), admixed FRO ancestry (green), local Ni-Vanuatu ancestry (dashed blue), Polynesian-related ancestry (dashed dark green), and intra-New Britain admixture (dotted blue). Colored bars give inferred total ancestry proportions (excluding outgroups Australian and Mixe). See Figure S5 for full results. S., southern.

the archipelago have experienced different demographic dynamics in the past.

The results in this study further our understanding of three population movements (M1–M3) that contributed substantially to the genetic makeup of Vanuatu through time, with new evidence presented pertaining to several open questions.

Four newly reported individuals from Teouma (Efate) join published data to make a total of 12 sampled Lapita-associated individuals (all represented by petrous bones) from Remote Oceania dating to 3000–2500 BP (eight from Teouma, three from Talasiu in Tonga, and one from Malakula), all of whom have nearly entirely FRO-related ancestry [4–6]. Thus, although future sampling could potentially still reveal greater genetic diversity during this period, ancient DNA results to date support the hypothesis that the first people of Remote Oceania, who were responsible for spreading the Lapita cultural complex (M1), were mostly descended from a population with roots in East and Southeast Asia [4].

After about 2500 BP, sampled individuals from post-Lapita contexts testify to an influx of Papuan ancestry (M2), although with different trajectories in different parts of Vanuatu. The three earliest individuals from this period from central and southern Vanuatu (one newly reported here) have the smallest proportions of FRO ancestry in our dataset, pointing to a major local genetic shift. The increased FRO ancestry in later populations from the same islands, combined with estimated dates of admixture that postdate the Lapita period, shows that mixture subsequently occurred between populations with different proportions of FRO and Papuan ancestry [5, 6]. Previously published late-Lapita and post-Lapita individuals (2500–2000 BP) from

Malakula in northern Vanuatu provide direct documentation of such an admixture process, as reflected in widely varying individual-level ancestry proportions along with recent estimated dates of admixture [5] (cf. Verdu and Rosenberg [25]). Unlike the other ancient individuals, those from Malakula come from a site that was continuously occupied for 1,000 years, from the founding Lapita population until around 2000 BP. There are also indications that elements of the Lapita culture persisted for longer in this region than in central and southern Vanuatu [28, 29].

Our reanalysis of ancient and present-day data supports a single source for the main component of Papuan ancestry found in Vanuatu from 2500 BP to the present, with most of the (few) exceptions potentially relating to post-European-contact movements. In particular, though we do not have contemporaneous ancient DNA data available from Near Oceania, the location of this source, based on the strong present-day regional genetic structure, is likely to have been New Britain, and we do not detect more than isolated evidence of gene flow from the (geographically closer) Solomon Islands (in agreement with Pugach et al. [11]). This relative homogeneity (across Vanuatu as well as through time) favors the hypothesis of a short-term migration episode responsible for introducing Papuan ancestry beginning around the late-Lapita period. Inferred dates of admixture in Vanuatu (aside from Polynesian-influenced groups) also point to mixture of FRO and Papuan ancestry around this time [5, 6]. *A priori*, the most likely movements and interactions would be expected to be between neighboring archipelagoes rather than distant ones, i.e., from the main Solomons chain to the Reefs and Santa Cruz to Vanuatu. However, this appears not to have

be the case either for M1, on archaeological and linguistic grounds [30], or for M2, on the basis of direct genetic links between Vanuatu and New Britain to the exclusion of the Solomons.

In light of results from both genetics and archaeology, a parsimonious explanation could be that M2 was effectively a continuation of M1 in late-Lapita times but involving migrants having mostly different ancestry. Cultural connections between New Britain and Vanuatu include the presence of New Britain obsidian in earliest Lapita deposits in Vanuatu [31], changes in dietary and mortuary behaviors and skeletal morphology subsequent to this earliest Lapita phase [32, 33], and distinctive practices (of unknown time depth), such as head binding and the production of fully circular pig's tusks, that are exclusive to those locations [5, 34]. We also find that, contrary to the more complex proposals in previous studies [5, 6], we can model the Papuan ancestry found in Polynesians by using the same New Britain-related source as for Vanuatu, raising the possibility that both were derived predominantly from the same phase of migration. However, as with the FRO component, future work is necessary to determine whether or not people carrying this ancestry passed through Vanuatu en route to Polynesia.

In accordance with archaeological and anthropological evidence of Polynesian cultural influence in Efate over the past several centuries, our analysis of five individuals from the Chief Roi Mata's Domain World Heritage Area demonstrates an influx of Polynesian-related ancestry as well (M3) through signals of higher FRO ancestry proportions, relatively recent dates of admixture, and specifically high allele sharing with Polynesians. The present-day Polynesian Outlier community of Mele, as well as other present-day and recent-past individuals from Efate and nearby islands (but not more distant groups), also display shared ancestry with the Eretok and Mangaas individuals, whereas the Polynesian Outlier population of Futuna and the neighboring island of Aneityum in southern Vanuatu likely represent a separate instance of Polynesian influence (we currently lack data for comparison from communities such as those of Lelepa and Mangaliliu in the immediate World Heritage Area vicinity). Thus, although the ancestry of present-day Ni-Vanuatu groups can largely be traced to the early human history of the archipelago, later migrations—in particular of Polynesians—have also contributed to the genetic diversity of Vanuatu today.

## STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **RESOURCE AVAILABILITY**
  - Lead Contact
  - Materials Availability
  - Data and Code Availability
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
  - Teouma
  - Mangaas
  - Eretok
  - Taplins
  - Banana Bay

## ● METHOD DETAILS

- Ancient DNA laboratory procedures
- Bioinformatic processing
- Uniparental haplogroups and authentication
- Radiocarbon Dates

## ● QUANTIFICATION AND STATISTICAL ANALYSIS

- Dataset construction
- PCA
- Formal modeling of admixture
- Dates of admixture
- $f_4$  regression analysis
- Admixture graph fitting

## SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.cub.2020.09.035>.

## ACKNOWLEDGMENTS

We thank Ann Marie Lawson, Fatma Zalzal, Jonas Oppenheimer, Kimberly Callan, Kristin Stewardson, Matthew Ferry, Megan Michel, Nasreen Broomandkhoshbacht, Nicole Adamski, Kendra Sirak, and Francesca Candilio for ancient DNA laboratory work; Swapnil Mallick and Matthew Mah for bioinformatics; Iñigo Olalde for help with kinship analysis; Rebecca Bernardos and Zhao Zhang for other data processing assistance; Douglas J. Kennett for help with radiocarbon dating; Johannes Krause, Kathrin Nägele, and Cosimo Posth for providing the EFE005 data; Graeme K. Ward for archaeological contributions; and Nick Patterson for helpful comments. We acknowledge the permission to sample at the Musée de l'Homme from the CRMD Management Committee, and we thank the Muséum national d'Histoire naturelle (Musée de l'Homme) for access to collections (Eretok and Mangaas) and Martin Friess for assistance in sample selection. We also gratefully acknowledge the interest and support of Ralph Regenvanu and of the Chiefs of Lelepa and Mangaliliu, to whom one of us (M.S.) presented results of this work in December 2019. D.R. was supported by the National Institutes of Health (NIGMS GM100233), the John Templeton Foundation (grant 61220), and the Paul Allen Foundation (Allen Discovery Center grant); D.R. is also an Investigator of the Howard Hughes Medical Institute.

## AUTHOR CONTRIBUTIONS

M.S., R.P., and D.R. supervised the study. M.S., F.V., S.B., R.S., W.Z., H.B., and R.P. provided samples and assembled archaeological and anthropological materials and information. F.P. performed radiocarbon dating analysis. R.M. served as a liaison for the project with other stakeholders. M.L. and D.R. analyzed genetic data. O.C. and N.R. performed and supervised ancient DNA laboratory work. M.L., M.S., and D.R. wrote the manuscript with input from all coauthors.

## DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: March 30, 2020

Revised: July 14, 2020

Accepted: September 10, 2020

Published: October 15, 2020

## REFERENCES

1. Green, R.C. (1991). Near and Remote Oceania: Disestablishing 'Melanesia' in culture history. In *Man and a Half: Essays in Pacific anthropology and ethnobiology in honour of Ralph Bulmer*, A. Pawley, ed. (The Polynesian Society), pp. 491–502.

2. Bedford, S., and Spriggs, M. (2008). Northern Vanuatu as a Pacific crossroads: The archaeology of discovery, interaction and the emergence of the 'ethnographic present'. *Asian Perspect.* 47, 95–120.
3. Bedford, S., and Spriggs, M. (2018). The archaeology of Vanuatu: 3000 years of history across islands of ash and coral. In *The Oxford Handbook of Prehistoric Oceania*, E. Cochrane, and T. Hunt, eds. (Oxford University Press), pp. 162–184.
4. Skoglund, P., Posth, C., Sirak, K., Spriggs, M., Valentin, F., Bedford, S., Clark, G.R., Reepmeyer, C., Petchey, F., Fernandes, D., et al. (2016). Genomic insights into the peopling of the Southwest Pacific. *Nature* 538, 510–513.
5. Posth, C., Nägele, K., Colleran, H., Valentin, F., Bedford, S., Kami, K.W., Shing, R., Buckley, H., Kinaston, R., Walworth, M., et al. (2018). Language continuity despite population replacement in Remote Oceania. *Nat. Ecol. Evol.* 2, 731–740.
6. Lipson, M., Skoglund, P., Spriggs, M., Valentin, F., Bedford, S., Shing, R., Buckley, H., Phillip, I., Ward, G.K., Mallick, S., et al. (2018). Population turnover in Remote Oceania shortly after initial settlement. *Curr. Biol.* 28, 1157–1165.e7.
7. Blust, R. (2008). Remote Melanesia: One history or two? An addendum to Donohue and Denham. *Oceanic Linguistics* 47, 445–459.
8. Bedford, S., Blust, R., Burley, D., Cox, M., Kirch, P., Matisoo-Smith, E., Naess, A., Pawley, A., Sand, C., and Sheppard, P. (2018). Ancient DNA and its contribution to understanding the human history of the Pacific Islands. *Archaeol. Ocean.* 53, 205–219.
9. Bedford, S., Spriggs, M., Buckley, H., Valentin, F., Regenvanu, R., and Abong, M. (2010). A cemetery of first settlement: Teouma, South Efate, Vanuatu/ Un cimetière de premier peuplement: le site de Teouma, sud d'Efate, Vanuatu. In *Lapita: Oceanic Ancestors/Lapita: Ancêtres Océaniens*, C. Sand, and S. Bedford, eds. (Musée de Quai Branly/Somogy), pp. 140–161.
10. Valentin, F., Bedford, S., Buckley, H., and Spriggs, M. (2010). Lapita burial practices: Evidence for complex body and bone treatment at the Teouma Cemetery, Vanuatu, Southwest Pacific. *J. Island Coast. Archaeol.* 5, 212–235.
11. Pugach, I., Duggan, A.T., Merriwether, D.A., Friedlaender, F.R., Friedlaender, J.S., and Stoneking, M. (2018). The gateway from Near into Remote Oceania: new insights from genome-wide data. *Mol. Biol. Evol.* 35, 871–886.
12. R. Feinberg, and R. Scaglion, eds. (2012). *Polynesian Outliers: The state of the art* (University of Pittsburgh Press).
13. Flexner, J., Bedford, S., and Valentin, F. (2019). Who was Polynesian? Who was Melanesian? Hybridity and ethnogenesis in the South Vanuatu Outliers. *J. Soc. Archaeol.* 3, 403–426.
14. Carson, M. (2012). Recent developments in prehistory: Perspectives on settlement, chronology, inter-community relations, and identity formation. In *Polynesian Outliers: The state of the art*, R. Feinberg, and R. Scaglion, eds. (University of Pittsburgh Press), pp. 27–48.
15. Zinger, W., Valentin, F., Flexner, J., Bedford, S., Déroit, F., and Grimaud-Hervé, D. (2020). How to explain Polynesian Outliers' heterogeneity? In *Networks and Monumentality in the Pacific*, Proceedings of the XVIII UISPP World Congress, A. Hermann, F. Valentin, C. Sand, and E. Nolet, eds. (Archaeopress), pp. 62–77.
16. Garanger, J. (1972). *Archéologie des Nouvelles Hébrides: Contribution à la connaissance des îles du Centre*. (Publications de la Société des Océanistes). <https://doi.org/10.4000/books.sdo.859>.
17. Espirat, J., Guiart, J., Lagrange, M.S., and Renaud, M. (1973). *Système des Titres Electifs ou Hérititaires dans les Nouvelles-Hébrides Centrales d'Efate aux Iles Shepherd*. (Museum National d'Histoire Naturelle).
18. Bedford, S., Spriggs, M., Wilson, M., and Regenvanu, R. (1998). The Australian National University-National Museum of Vanuatu Archaeological Project 1994-7: A Preliminary Report on the Establishment of Cultural Sequences and Rock Art Research. *Asian Perspect.* 37, 165–193.
19. Petchey, F., Spriggs, M., Bedford, S., Valentin, F., and Buckley, H. (2014). Direct radiocarbon dating of burials from the Teouma Lapita cemetery, Efate, Vanuatu. *Journal of Archaeological Science* 50, 227–242.
20. Haak, W., Lazaridis, I., Patterson, N., Rohland, N., Mallick, S., Llamas, B., Brandt, G., Nordenfelt, S., Harney, E., Stewardson, K., et al. (2015). Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* 522, 207–211.
21. Lazaridis, I., Mitnik, A., Patterson, N., Mallick, S., Rohland, N., Pfrengle, S., Furtwängler, A., Peltzer, A., Posth, C., Vasilakis, A., et al. (2017). Genetic origins of the Minoans and Mycenaeans. *Nature* 548, 214–218.
22. Pickrell, J.K., Patterson, N., Loh, P.R., Lipson, M., Berger, B., Stoneking, M., Pakendorf, B., and Reich, D. (2014). Ancient west Eurasian ancestry in southern and eastern Africa. *Proc. Natl. Acad. Sci. USA* 111, 2632–2637.
23. Narasimhan, V.M., Patterson, N., Moorjani, P., Rohland, N., Bernardos, R., Mallick, S., Lazaridis, I., Nakatsuka, N., Olalde, I., Lipson, M., et al. (2019). The formation of human populations in South and Central Asia. *Science* 365, eaat7487.
24. Spriggs, M. (1997). *The Island Melanesians* (Blackwell Publishers).
25. Verdu, P., and Rosenberg, N.A. (2011). A general mechanistic model for admixture histories of hybrid populations. *Genetics* 189, 1413–1426.
26. Shineberg, D. (1967). *They Came for Sandalwood: A study of the sandalwood trade in the South West Pacific* (Melbourne University Press).
27. Reich, D., Patterson, N., Campbell, D., Tandon, A., Mazieres, S., Ray, N., Parra, M.V., Rojas, W., Duque, C., Mesa, N., et al. (2012). Reconstructing Native American population history. *Nature* 488, 370–374.
28. Bedford, S., Buckley, H., Valentin, F., Tayles, N., and Longga, N. (2011). Lapita burials, a new Lapita cemetery and Post-Lapita burials from Malakula, Northern Vanuatu, Southwest Pacific. *Journal of Pacific Archaeology* 2, 26–48.
29. Bedford, S. (2019). Lapita pottery from the small islands of northeast Malakula, Vanuatu: A brief overview and implications. In *Debating Lapita: Distribution, chronology, society and subsistence*, S. Bedford, and M. Spriggs, eds. (ANU Press), pp. 225–240.
30. Sheppard, P. (2019). Early Lapita colonisation of Remote Oceania: An update on the leapfrog hypothesis. In *Debating Lapita: Distribution, chronology, society and subsistence*, S. Bedford, and M. Spriggs, eds. (ANU Press), pp. 135–153.
31. Reepmeyer, C., Spriggs, M., Bedford, S., and Ambrose, W. (2011). Provenance and technology of lithic artefacts from the Teouma Lapita Site, Vanuatu. *Asian Perspect.* 49, 205–225.
32. Valentin, F., Herrscher, E., Bedford, S., Spriggs, M., and Buckley, H. (2014). Evidence for social and cultural change in Central Vanuatu during the first millennium BC: comparing funerary and dietary patterns of the first and later generations at Teouma, Efate. *J. Island Coast. Archaeol.* 9, 381–399.
33. Valentin, F., Déroit, F., Spriggs, M.J., and Bedford, S. (2016). Early Lapita skeletons from Vanuatu show Polynesian craniofacial shape: Implications for Remote Oceanic settlement and Lapita origins. *Proc. Natl. Acad. Sci. USA* 113, 292–297.
34. Bedford, S. (2018). Modified canines: Circular pig's tusks in Vanuatu and the wider Pacific. In *The Archaeology of Portable Art: Southeast Asian, Pacific, and Australian Perspectives*, M.C. Langley, M. Litster, D. Wright, and S.K. May, eds. (Routledge), pp. 125–141.
35. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.; 1000 Genome Project Data Processing Subgroup (2009). The sequence alignment/map format and SAMtools. *Bioinformatics* 25, 2078–2079.
36. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25, 1754–1760.
37. Patterson, N., Moorjani, P., Luo, Y., Mallick, S., Rohland, N., Zhan, Y., Genschorek, T., Webster, T., and Reich, D. (2012). Ancient admixture in human history. *Genetics* 192, 1065–1093.

38. Patterson, N., Price, A.L., and Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genet.* 2, e190.
39. Skoglund, P., Northoff, B.H., Shunkov, M.V., Derevianko, A.P., Pääbo, S., Krause, J., and Jakobsson, M. (2014). Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. USA* 111, 2229–2234.
40. Weissensteiner, H., Pacher, D., Kloss-Brandstätter, A., Forer, L., Specht, G., Bandelt, H.J., Kronenberg, F., Salas, A., and Schönherr, S. (2016). HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res.* 44 (W1), W58–63.
41. Fu, Q., Mittnik, A., Johnson, P.L.F., Bos, K., Lari, M., Bollongino, R., Sun, C., Gienschi, L., Schmitz, R., Burger, J., et al. (2013). A revised timescale for human evolution based on ancient mitochondrial genomes. *Curr. Biol.* 23, 553–559.
42. Bronk Ramsey, C. (1995). Radiocarbon calibration and analysis of stratigraphy: The OxCal program. *Radiocarbon* 37, 425–430.
43. Valentin, F., Allié, F., Bedford, S., and Spriggs, M. (2016). Réflexions sur la transformation anthropique du cadavre: Le cas des sépultures Lapita de Teouma (Vanuatu). *Bull. Mem. Soc. Anthropol. Paris* 28, 39–44.
44. Bedford, S. (2006). Pieces of the Vanuatu Puzzle: Archaeology of the North, South and Centre. *Terra Australis* 23.
45. Spriggs, M., and Bedford, S. (2001). Arapus: A Lapita site at Mangaasi in central Vanuatu? In *The archaeology of Lapita dispersal in Oceania*, G.R. Clark, A.J. Anderson, and T. Vunidilo, eds. (Pandanus Books), pp. 93–104.
46. Ward, G., and Houghton, P. (1991). The Mele burials (Vanuatu): salvage excavations and biological relationships. *Indo-Pacific Prehistory Association Bulletin* 11, 229–235.
47. Valentin, F., Spriggs, M., Bedford, S., and Buckley, H. (2011). Vanuatu mortuary practices over three millennia: Lapita to the Early European Contact Period. *Journal of Pacific Archaeology* 2, 49–65.
48. Shing, R., and Philip, I. (2010). Preliminary report on the burial 1, Banana Bay. Report on file at Vanuatu Cultural Centre.
49. Dabney, J., Knapp, M., Glocke, I., Gansauge, M.T., Weihmann, A., Nickel, B., Valdiosera, C., Garcia, N., Pääbo, S., Arsuaga, J.L., and Meyer, M. (2013). Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. USA* 110, 15758–15763.
50. Korlević, P., Gerber, T., Gansauge, M.T., Hajdinjak, M., Nagel, S., Aximu-Petri, A., and Meyer, M. (2015). Reducing microbial and human contamination in DNA extractions from ancient bones and teeth. *Biotechniques* 59, 87–93.
51. Rohland, N., Glocke, I., Aximu-Petri, A., and Meyer, M. (2018). Extraction of highly degraded DNA from ancient bones, teeth and sediments for high-throughput sequencing. *Nat. Protoc.* 13, 2447–2461.
52. Sirak, K.A., Fernandes, D.M., Cheronet, O., Novak, M., Gamarra, B., Balassa, T., Bernert, Z., Cséki, A., Dani, J., Gallina, J.Z., et al. (2017). A minimally-invasive method for sampling human petrous bones from the cranial base for ancient DNA analysis. *Biotechniques* 62, 283–289.
53. Briggs, A.W., Stenzel, U., Meyer, M., Krause, J., Kircher, M., and Pääbo, S. (2010). Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res.* 38, e87.
54. Meyer, M., and Kircher, M. (2010). Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* 2010, t5448.
55. Rohland, N., Harney, E., Mallick, S., Nordenfelt, S., and Reich, D. (2015). Partial uracil-DNA-glycosylase treatment for screening of ancient DNA. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 370, 20130624.
56. Gansauge, M.-T., Aximu-Petri, A., Nagel, S., and Meyer, M. (2020). Manual and automated preparation of single-stranded DNA libraries for the sequencing of DNA from ancient biological remains and other sources of highly degraded DNA. *Nat. Protoc.* 15, 2279–2300.
57. Fu, Q., Meyer, M., Gao, X., Stenzel, U., Burbano, H.A., Kelso, J., and Pääbo, S. (2013). DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl. Acad. Sci. USA* 110, 2223–2227.
58. Fu, Q., Hajdinjak, M., Moldovan, O.T., Constantin, S., Mallick, S., Skoglund, P., Patterson, N., Rohland, N., Lazaridis, I., Nickel, B., et al. (2015). An early modern human from Romania with a recent Neanderthal ancestor. *Nature* 524, 216–219.
59. Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S.A., Harney, E., Stewardson, K., Fernandes, D., Novak, M., et al. (2015). Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* 528, 499–503.
60. Lazaridis, I., Nadel, D., Rollefson, G., Merrett, D.C., Rohland, N., Mallick, S., Fernandes, D., Novak, M., Gamarra, B., Sirak, K., et al. (2016). Genomic insights into the origin of farming in the ancient Near East. *Nature* 536, 419–424.
61. Kircher, M., Sawyer, S., and Meyer, M. (2012). Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res.* 40, e3.
62. Behar, D.M., van Oven, M., Rosset, S., Metspalu, M., Loogväli, E.L., Silva, N.M., Kivisild, T., Torroni, A., and Villems, R. (2012). A “Copernican” reassessment of the human mitochondrial DNA tree from its root. *Am. J. Hum. Genet.* 90, 675–684.
63. Skoglund, P., Storå, J., Götherström, A., and Jakobsson, M. (2013). Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* 40, 4477–4482.
64. Korneliusson, T.S., Albrechtsen, A., and Nielsen, R. (2014). ANGSD: Analysis of next generation sequencing data. *BMC Bioinformatics* 15, 356.
65. Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., et al. (2013). IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 Years cal BP. *Radiocarbon* 55, 1869–1887.
66. Petchey, F., Spriggs, M., Leach, F., Seed, M., Sand, C., Pietruszewski, M., and Anderson, K. (2011). Testing the human factor: Radiocarbon dating the first peoples of the South Pacific. *J. Archaeol. Sci.* 38, 29–44.
67. Petchey, F., Anderson, A., Zondervan, A., Ulm, S., and Hogg, A. (2008). New marine  $\Delta R$  values for the South Pacific subtropical gyre region. *Radiocarbon* 50, 373–397.
68. Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O., Marchini, J.L., McCarthy, S., McVean, G.A., and Abecasis, G.R.; 1000 Genomes Project Consortium (2015). A global reference for human genetic variation. *Nature* 526, 68–74.
69. Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M., Chennagiri, N., Nordenfelt, S., Tandon, A., et al. (2016). The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* 538, 201–206.
70. Loh, P.R., Lipson, M., Patterson, N., Moorjani, P., Pickrell, J.K., Reich, D., and Berger, B. (2013). Inferring admixture histories of human populations using linkage disequilibrium. *Genetics* 193, 1233–1254.

## STAR★METHODS

### KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Biological Samples		
Ancient human skeletal elements	This study	See <a href="#">Table 1</a>
Chemicals, Peptides, and Recombinant Proteins		
Pfu Turbo Cx Hotstart DNA Polymerase	Agilent Technologies	600412
Herculase II Fusion DNA Polymerase	Agilent Technologies	600679
2x HI-RPM hybridization buffer	Agilent Technologies	5190-0403
0.5 M EDTA pH 8.0	BioExpress	E177
Silica magnetic beads	G-Biosciences	786-915
Sera-Mag™ Magnetic Speed-beads™ Carboxylate-Modified (1 μm, 3EDAC/PA5)	GE LifeScience	65152105050250
USER enzyme	New England Biolabs	M5505
UGI	New England Biolabs	M0281
Bst DNA Polymerase2.0, large frag.	New England Biolabs	M0537
PE buffer concentrate	QIAGEN	19065
Proteinase K	Sigma Aldrich	P6556
Guanidine hydrochloride	Sigma Aldrich	G3272
3M Sodium Acetate (pH 5.2)	Sigma Aldrich	S7899
Water	Sigma Aldrich	W4502
Tween-20	Sigma Aldrich	P9416
Isopropanol	Sigma Aldrich	650447
Ethanol	Sigma Aldrich	E7023
5M NaCl	Sigma Aldrich	S5150
1M NaOH	Sigma Aldrich	71463
20% SDS	Sigma Aldrich	5030
PEG-8000	Sigma Aldrich	89510
1 M Tris-HCl pH 8.0	Sigma Aldrich	AM9856
dNTP Mix	Thermo Fisher Scientific	R1121
ATP	Thermo Fisher Scientific	R0441
10x Buffer Tango	Thermo Fisher Scientific	BY5
T4 Polynucleotide Kinase	Thermo Fisher Scientific	EK0032
T4 DNA Polymerase	Thermo Fisher Scientific	EP0062
T4 DNA Ligase	Thermo Fisher Scientific	EL0011
Maxima SYBR Green kit	Thermo Fisher Scientific	K0251
50x Denhardt's solution	Thermo Fisher Scientific	750018
SSC Buffer (20x)	Thermo Fisher Scientific	AM9770
GeneAmp 10x PCR Gold Buffer	Thermo Fisher Scientific	4379874
Dynabeads MyOne Streptavidin T1	Thermo Fisher Scientific	65602
Salmon sperm DNA	Thermo Fisher Scientific	15632-011
Human Cot-I DNA	Thermo Fisher Scientific	15279011
DyNAmo HS SYBR Green qPCR Kit	Thermo Fisher Scientific	F410L
Methanol, certified ACS	VWR	EM-MX0485-3
Acetone, certified ACS	VWR	BDH1101-4LP
Dichloromethane, certified ACS	VWR	EMD-DX0835-3
Hydrochloric acid, 6N, 0.5N & 0.01N	VWR	EMD-HX0603-3

(Continued on next page)



**Continued**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Critical Commercial Assays		
High Pure Extender from Viral Nucleic Acid Large Volume Kit	Roche	5114403001
NextSeq® 500/550 High Output Kit v2 (150 cycles)	Illumina	FC-404-2002
Deposited Data		
Raw and analyzed data	This paper	ENA: PRJEB40109
Software and Algorithms		
In-house bioinformatics tools	<a href="https://github.com/DReichLab/ADNA-Tools">https://github.com/DReichLab/ADNA-Tools</a>	N/A
In-house data workflow	<a href="https://github.com/DReichLab/adna-workflow">https://github.com/DReichLab/adna-workflow</a>	N/A
Samtools	[35]	<a href="http://samtools.sourceforge.net/">http://samtools.sourceforge.net/</a>
BWA	[36]	<a href="http://bio-bwa.sourceforge.net/">http://bio-bwa.sourceforge.net/</a>
Picard	<a href="https://broadinstitute.github.io/picard/">https://broadinstitute.github.io/picard/</a>	N/A
ADMIXTOOLS	[37]	<a href="https://github.com/DReichLab/AdmixTools">https://github.com/DReichLab/AdmixTools</a>
SeqPrep	<a href="https://github.com/jstjohn/SeqPrep">https://github.com/jstjohn/SeqPrep</a>	N/A
bamrmdup	<a href="https://bitbucket.org/ustenzel/biohazard">https://bitbucket.org/ustenzel/biohazard</a>	N/A
smartpca	[38]	<a href="https://www.hsph.harvard.edu/alkes-price/software/">https://www.hsph.harvard.edu/alkes-price/software/</a>
PMDtools	[39]	<a href="https://github.com/pontusssk/PMDtools">https://github.com/pontusssk/PMDtools</a>
Haplogrep 2	[40]	<a href="http://haplogrep.uibk.ac.at/">http://haplogrep.uibk.ac.at/</a>
htsbox	<a href="https://github.com/lh3/htsbox">https://github.com/lh3/htsbox</a>	N/A
contamMix	[41]	contact Philip Johnson <a href="mailto:plfj@umd.edu">plfj@umd.edu</a>
OxCal	[42]	<a href="https://c14.arch.ox.ac.uk/oxcal.html">https://c14.arch.ox.ac.uk/oxcal.html</a>
MALDER	[22]	<a href="https://github.com/joepickrell/malder/tree/master/MALDER">https://github.com/joepickrell/malder/tree/master/MALDER</a>
DATES	[23]	<a href="https://github.com/priyamoorejani/DATES">https://github.com/priyamoorejani/DATES</a>

**RESOURCE AVAILABILITY****Lead Contact**

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, David Reich ([reich@genetics.med.harvard.edu](mailto:reich@genetics.med.harvard.edu)).

**Materials Availability**

This study did not generate new unique reagents.

**Data and Code Availability**

The aligned sequences are available through the European Nucleotide Archive under accession number ENA: PRJEB40109. Genotype data files are available at <https://reich.hms.harvard.edu/datasets>.

**EXPERIMENTAL MODEL AND SUBJECT DETAILS**

The following are brief descriptions of the sites and burials where the ancient individuals included in this study were found.

**Teouma**

The Teouma site is located in southern Efate, on the edge of a large sheltered bay. It was once located near the sea but due to uplift it is now some 800 m from the current coast and 8 m above sea level. It comprises a colonizing Lapita settlement and associated cemetery dating from c. 2950 BP. Six field seasons of excavation were carried out at the site [9]. They revealed an extensive cemetery with up to 68 burial features. The burials were placed in solution holes in the ancient uplifted reef or in shallow graves on the old coral beach foreshore. They were directly associated with Lapita pottery and a range of ornaments also typical of Lapita. Manipulation

of the bodies and the skeletal remains was standard procedure with all the adult skulls being removed from the initial interments [10, 43]. In a number of rare cases skulls were placed within other graves. Petrous portions of four of them were analyzed here (B10A, B10C, B30B, B30C). Continuing Lapita-period occupation at the site along with subsequent post-Lapita occupation ultimately buried the cemetery. The site appears to have been abandoned by about 2500 BP.

### Mangaas

The Mangaas or Mangaasi site is located on the west coast of Efate opposite Lelepa Island. The site was first excavated by José Garanger in 1967 as part of his wider pioneering archaeological research on central Vanuatu [16]. In oral traditions the site is said to be the location of the village of Roi Mata, a powerful chief who transformed the socio-political organization of the region. Deeply buried deposits were identified that were associated with distinctive pottery, subsequently named Mangaasi. Two burials (represented by petrous bones) recovered in the upper layers of the site are the subject of analysis here while five graves and two groups of disturbed human remains were recorded by Garanger. The same site and a much more extensive area immediately adjacent to the west were subsequently re-investigated from 1996–2003 [44, 45]. It has now been established that the region was first occupied around 2800 BP with continuing settlement in the region up to the present, primarily focused parallel to the coast. The earliest settlement is now some 80 m from the current beach due to continuing uplift, and, over millennia, settlements have continued to shift to maintain their location near the coast. The earlier archaeological deposits are generally deeply buried due to subsequent slopewash and tephra deposits.

### Eretok

Eretok (also known as Retoka or Hat Island) is located just offshore of Efate and Lelepa Islands on the west coast of Efate. It is the location of a cemetery that was associated with the death of chief Roi Mata in c. 1600 CE. Oral traditions tell of the death of this very important chief and how subsequently he was buried as part of a large communal ceremony undertaken on the island. Dozens of people apparently volunteered to be buried with the chief as part of the ceremony. The site was excavated by José Garanger in 1967 after he was informed of its location by local community members working on the site of Roi Mata's village at Mangaas [16]. More than 50 individuals were identified with many buried as couples and others individually. Three of them, represented by two petrous bones and one tooth, are analyzed here. Roi Mata is identified as being buried in a more deeply excavated zone in front of a series of standing stones, alongside a number of individuals ostentatiously decorated with traditional shell and other ornaments.

### Taplins

Taplins comprises two rockshelters, Taplins 1 and 2, located at the base of a cliff on uplifted terraces behind Mele Bay in the south-western part of Efate. Five subsurface graves were excavated at these sites by Graeme Ward in 1973 and 1974 [46, 47]. Both the earlier analyzed individual and the subject of this study came from Taplins 1. The loose tooth studied here was initially hypothesized to belong to the same individual as the previously published petrous bone sample [6], but the genetic analysis shows that a second individual is represented (different mtDNA and Y chromosome haplogroups, and genome-wide allele-matching rates at the level of unrelated individuals).

### Banana Bay

Four burials were located during drainage works associated with road improvements around Efate Island [48]. The site is located on the southeast coast of the island. Local informants said that there had been a large village located in this area up to European contact. Burial 1, a burial in a supine position some 1.5 m below the current ground surface, was clearly associated with the historic period as the body was adorned with a shell and glass bead necklace. That individual was analyzed in ref. [6]. The tooth studied here is associated with a group of bones representing at least one other individual, found close by burial 1.

## METHOD DETAILS

### Ancient DNA laboratory procedures

For the Teouma and Taplins samples, powder was drilled from bones or teeth in a clean room facility at University College Dublin, and DNA was then extracted in dedicated clean rooms at Harvard Medical School following previously published protocols [49–51] (additional sample preparation information can be found in Data S1A). Powder was obtained from four of the Mangaas and Eretok samples via cranial base drilling [52] at the Musée de l'Homme in Paris, while for I14493, the drilling step was omitted, and the tooth was submerged directly in 1.5 mL of extraction buffer for 4 h. Laboratory work for EFE005 took place at the Max Planck Institute for the Science of Human History in Jena, Germany. The tooth was cut along the enamel/dentin junction and drilled into the pulp chamber, with the extraction then proceeding as above.

Barcoded sequencing libraries (1–5 per individual) were prepared from the extracts, utilizing the enzyme uracil-DNA glycosylase (UDG; partial treatment, for all but EFE005) to reduce the rate of deamination-induced ancient DNA damage artifacts [53–56]. The libraries were enriched for sequence fragments overlapping the mitochondrial genome and ~1.2 million genome-wide SNPs via two rounds of in-solution target capture [20, 57–60], with 7-base-pair indices added for the libraries generated at Harvard Medical School [61]. The libraries were sequenced on an Illumina HiSeq machine with single-end reads (EFE005) or an Illumina NextSeq 500 machine with 76-base paired-end reads (others).

### Bioinformatic processing

For the ten individuals for whom data were generated at Harvard Medical School, we assigned sequencing reads to their respective libraries based on their barcodes, requiring at most one mismatch per read pair. We merged overlapping reads, trimmed barcodes and adapters, and then mapped to the mitochondrial reference genome RSRS [62] and to the human reference genome (version hg19) using the ‘samse’ command with default parameters in BWA (version 0.6.1) [36]. After aligning, we removed duplicate molecules and imposed a mapping quality filter of 10. Finally, we trimmed terminal bases (2 for UDG-treated libraries and 5 for untreated) to eliminate most damage-induced errors, and we called pseudo-haploid genotypes for genome-wide analyses by selecting one allele at random per targeted SNP site. Data for EFE005 were processed at the Max Planck Institute in Jena as described elsewhere [5].

### Uniparental haplogroups and authentication

We determined genetic sex of each individual by examining the fractions of sequence fragments mapping to the X and Y chromosomes [63]. We called mitochondrial haplogroups using HaploGrep2 [40] and Y chromosome haplogroups by comparing SNP genotypes (using all reads) to the International Society of Genetic Genealogy Y-tree (<http://www.isogg.org>).

We assessed the authenticity of the data through five measures (Data S1A). First, we computed the rate of damage-induced errors in terminal positions of sequenced molecules to confirm the presence of ancient DNA signatures. We then tested for possible contamination by (a) confirming that genetic sex could be determined as male or female, (b) computing the rate of matching of mtDNA sequences to the consensus haplogroup call for each individual [41], and (c) measuring apparent heterozygosity at variable sites on the X chromosome in males [64]. Finally, we noted any signals in the genome-wide ancestry analyses that could suggest possible contamination from present-day human DNA.

For the individuals with lower coverage (fewer than 100,000 SNPs), the metrics are noisier, and the contamination estimates are generally less reliable, so our typical approach was to run our analyses for these individuals but to be cautious in interpreting the results and not to draw fine-grained conclusions. For the higher-coverage individuals, all metrics indicated at most a few percent contamination. One individual (I14493) had lower than expected damage rates (2.4% for mapped nuclear reads) but low contamination estimates (about 2%–5% from both mtDNA and X chromosome). As an empirical test, we fit an admixture model for I14493 in *qpAdm* using damage-restricted data, and the results were extremely similar to those for all data ( $p = 0.61$ ,  $82.7 \pm 4.4\%$  Baining-related ancestry, and  $17.3 \pm 4.4\%$  Kankanaey-related ancestry, versus  $p = 0.84$ ,  $79.8 \pm 1.4\%$ , and  $20.2 \pm 1.4\%$ ; Data S1D). Thus, we continued to use the data in our analyses.

### Radiocarbon Dates

We report new direct AMS radiocarbon dates for three individuals (EFE005, I10967, and I14493), which we combined with previously published dates for I5267 and I5268 [19]. Dates were calibrated using OxCal [42] version 4.3 with a mixture of the Marine13 and Intcal13 curves [65] as determined by linear interpolation between dietary terrestrial and marine  $\delta^{13}\text{C}$  isotopic endpoints ( $-21\text{‰}$ – $-12\text{‰}$ ) with an uncertainty of  $\pm 10\%$  on the percent marine carbon result, following the methodology outlined in ref. [66] to assess the proportion of marine, reef, and terrestrial food contribution to the bone protein. A location-specific reservoir correction (DR) of  $40 \pm 44$   $^{14}\text{C}$  years was also applied to the marine curve to adjust for regional oceanic variation in  $^{14}\text{C}$  around Vanuatu [67].

## QUANTIFICATION AND STATISTICAL ANALYSIS

### Dataset construction

We merged our newly generated data with published ancient and present-day data [4–6, 68, 69]. Unless otherwise specified, we used a set of ~398,000 autosomal SNPs from the Human Origins array, excluding (a) C-to-T transition SNPs at CpG dinucleotides, and (b) a set of SNPs with high rates of missing data in present-day genotype data. For *f*-statistic-based analyses (as reflected in the sample sizes in Data S1B), we excluded 20 present-day individuals who were outliers relative to their ethnolinguistic groups: UV128 (Tolai); UV219 (Mengen); UV220 (Sulka); UV726 (Kuot Kabil); UV516 and UV519 (Kuot Lamalau); UV533 (Nailik); UV1166 (Melamela); Jk2663, Jk2665, and Jk2669 (Samoan); and nine individuals from Vanuatu who were identified as outliers in previously published data curations [6].

### PCA

We performed PCA with smartpca [38], using the ‘lsqproject’ and ‘shrinkmode’ options. We used three populations (Kankanaey, Nasioi, and New Guinea Highlanders) to define axes and projected all other individuals. Projecting ancient individuals prevents bias due to missing data; we chose to project present-day populations as well in order to create a two-dimensional plot with equivalent procedures for all individuals (aside from the three axis populations) and with minimal effects of population-specific drift. We note that the ancient individuals with lower coverage have more uncertainty associated with their positions.

### Formal modeling of admixture

We tested admixture models using the *qpAdm* software [20, 21]. Our basic outgroup list consisted of New Guinea, Australian, French, Dai, Onge, and Mixe, a set of populations with largely phylogenetically distinct positions relative to the mixing populations in our applications: Papuan, deeply Papuan-related, western Eurasian, East Asian, deep eastern Eurasian, and Native American, respectively.

If a given model fits poorly (i.e., low  $p$  value), that implies that it is poorly specified, in the sense that not all of the ancestry in the test population is more closely related to the proxy sources than to the outgroups. In other words, either the test population shares some ancestry with one or more of the outgroups more closely than with the specified proxy sources, or one of the proxy sources shares ancestry with one or more of the outgroups more closely than with the test population. To search for possible sex-biased admixture, we compared mixture proportion estimates on the autosomes and the X chromosome, computing a quasi-Z-score by dividing the difference by the standard error for the X estimate (which is much larger than the autosomal standard error). To maximize coverage on the X chromosome, we did not apply the two SNP exclusion criteria described in the ‘Dataset construction’ subsection above (which should have a negligible effect on mixture proportion estimates in the units of ratios of  $f$ -statistics). When computing FRO and Papuan ancestry proportions, we used Baining as the proxy for Papuan-related ancestry and corrected the estimates for the fact that Baining themselves have ~5% FRO ancestry.

We also tested the compatibility of multiple populations with having common sources of admixture without a formal model, using *qpWave* [27]. Our test set for ancient Vanuatu plus other Oceanian populations consisted of present-day Tongan (6 individuals) and Melamela (9), plus the following ancient Vanuatu groupings: Efate 150–400 BP (5), Efate ~600 BP (1), Efate ~2400 BP (2), Epi ~150 BP (2), Epi ~1300 BP (2), Futuna ~1100 BP (4), Malakula 2000–2500 BP (6), Tanna ~150 and ~2500 BP (2), Mangaas (2), and Eretok (3). For present-day Vanuatu, we used all population groups in our dataset. In both analyses, our outgroup set was the same as for *qpAdm*, either with or without Nasioi. As in *qpAdm*, a higher  $p$  value indicates a better fit for the proposed model, where a rank of  $k$  implies  $k+1$  distinct ancestry sources combining to form the test set of populations.

### Dates of admixture

We estimated dates of admixture using MALDER [22] and DATES [23]. MALDER extends the linkage disequilibrium (LD)-based model of ALDER [70] by integrating information from multiple reference populations and searching for evidence of multiple waves of admixture. We used all ~590k autosomal Human Origins SNPs, and our reference set consisted of New Guinea Highlanders, Papuan, Australian, Baining (both subgroups), Teouma, Talasiu, Kankanaey, Ami, and CDX (1000 Genomes Dai).

DATES implements a regression-based ancestry covariance estimate that can be applied to single individuals. We used all ~1.15 million autosomal SNPs from our capture set, and our reference pair was Papuan [69] and CDX [68]. For both methods, we assumed an average generation interval of 28 years when converting results to years in the past and estimated standard errors by block jackknife.

### $f_4$ regression analysis

We used a linear regression-based method to test for asymmetrical allele-sharing in cases where the  $f_4$ -statistics of interest are confounded by differential ancestry proportions across the test population set. Instead of searching directly for non-zero values, we plotted pairs of  $f_4$ -statistics in which the dependent variable is the statistic of interest and the independent variable is a statistic ( $f_4(X, \text{New Guinea Highlanders; Kankanaey, Australian})$ ) measuring levels of Papuan ancestry. This approach is based on the linearity of  $f_4$ -statistics for a collection of test populations (‘X’) with mixtures of ancestry related to the same two source populations but in different proportions (see below for derivation). If some test populations violate the proposed two-way model, they will tend to deviate from the expected linear relationship between the dependent and independent variables. We computed the  $f_4$ -statistics in ADMIX-TOOLS [37], with standard errors estimated by block jackknife. This approach is in some ways similar to the  $f_4$ -biplots introduced in ref. [20], but the scenarios of interest and the interpretations of the results are different.

Taking the example of Figure 4B, suppose we had a collection of populations, each of whose ancestry is a mixture from the same two sources, P and F, but in different proportions. Let  $A = f_4(P, \text{New Guinea Highlanders; Kankanaey, Australian})$ ,  $B = f_4(F, \text{New Guinea Highlanders; Kankanaey, Australian})$ ,  $C = f_4(P, \text{Tolai; Kankanaey, Tongan})$ , and  $D = f_4(F, \text{Tolai; Kankanaey, Tongan})$ . If one of our test populations, X, has a proportion  $a$  of P-related ancestry and  $(1-a)$  of F-related ancestry, then (in expectation)  $f_4(X, \text{New Guinea Highlanders; Kankanaey, Australian}) = aA + (1-a)B = a(A-B) + B$ , and  $f_4(X, \text{Tolai; Kankanaey, Tongan}) = aC + (1-a)D = a(C-D) + D = [\text{constant1}] * f_4(X, \text{New Guinea Highlanders; Kankanaey, Australian}) + [\text{constant2}]$  (where the first constant is  $(C-D)/(A-B)$  and the second is also a rational function of A, B, C, and D). Thus a pair of  $f_4$ -statistics are expected to have a linear relationship under the assumption that the set of populations in the first position (with the other three positions fixed) have mixtures of ancestry from the same two sources.

We performed linear regression via inverse-variance-weighted least-squares. Given the resulting best-fit equation  $f_4(2) = m * f_4(1) + b$ , we evaluated the deviation of each population by calculating its empirical value of  $f_4(2) - m * f_4(1) - b$ , assessing the statistical significance by a Z-test (estimating the standard error on the value directly with a block jackknife). In most cases, we used one data point for each population group, except in cases of ancient populations with substantial heterogeneity (the distinction being accommodated naturally because of the weighting scheme).

To maximize power given the relatively low-coverage data for the Lapita-period individuals, we computed the statistic  $f_4(X, \text{New Guinea Highlanders; Teouma, Talasiu})$  indirectly, via  $f_4(X, \text{New Guinea Highlanders; CDX, Talasiu}) - f_4(X, \text{New Guinea Highlanders; CDX, Teouma})$  (including non-overlapping SNPs), with a block jackknife to estimate the standard error. When computing deviations from the regression line for this statistic, we then used the raw standard error rather than the full residual described above; empirically, this likely results in slight underestimates of the standard error (although this is conservative, in the sense that we observe only minor deviations for this test).

We note that the choice of comparison population in the second position (e.g., New Guinea in the previous paragraph) only serves to shift all statistic values up or down a constant amount, because  $f_4(X, \text{Pop1}; Y, Z) - f_4(X, \text{Pop2}; Y, Z) = f_4(\text{Pop2}, \text{Pop1}; Y, Z)$ , which is a constant for all  $X$ . For Polynesian-related ancestry tests, to improve power, we used Tolai in the second position because (a) it is the Oceanian population with the largest sample size in our dataset, (b) it has an intermediate proportion of FRO ancestry, and (c) we were not specifically interested in the history of Tolai as a test population in these analyses.

When testing for specific relatedness to the Eretok and Mangaas individuals, we used the statistics  $f_4(X, \text{Tolai}; \text{Eretok/Mangaas individual}, \text{Futuna} \sim 1100 \text{ BP})$  for each individual in turn. We used the set of four ancient individuals from Futuna [5] in the fourth position rather than a present-day group in order to prevent artificial signals of allele-sharing when  $X$  is ancient.

### Admixture graph fitting

We built our admixture graph using the *qpGraph* software in ADMIXTOOLS [37], with 13 populations included: Mixe (from Mexico) and Australian as outgroups; Atayal and Kankanaey (FRO-related); New Guinea Highlanders; Nasioi (Solomon Islands); Baining (Marabu subgroup) and Melamela (New Britain); Tongan; and four groups or individuals from Vanuatu – Eretok and Mangaas, ~600 BP Efate (15259) [6], present-day Futuna, and present-day Tanna. We specified the options ‘outpop: NULL’, ‘lambdascale: 1’, and ‘diag: 0.0001.’ For a model of this size, the space of possible topologies is extremely large, so we cannot conclude that our final graph is the unique one that provides a good fit to the data. Instead, we use it in conjunction with our other analyses to investigate which results are supported when modeling the relationships among many populations simultaneously and to discover any additional admixture events necessary to obtain a good fit.